

Available online at www.sciencerepository.org

Science Repository



Research Article

Extracellular Matrix Expression in Human Induced Pluripotent Stem Cell-Derived Optic Vesicles

Ramesh R. Kaini², Christina L. Rettinger² and Heuy-Ching Wang^{1,2*}

¹Combat Casualty Care and Operational Medicine, Naval Medical Research Unit-San Antonio (NAMRU-SA), Fort Sam Houston, San Antonio, Texas, USA

²Ocular and Sensory Trauma Task Area, United States Army Institute of Surgical Research, Fort Sam Houston, San Antonio, Texas, USA

ARTICLE INFO

Article history:

Received: 16 August, 2021

Accepted: 2 September, 2021

Published: 21 September, 2021

Keywords:

extracellular matrix

induced pluripotent stem cells

optic vesicles

retina

embryoid body

retinogenesis

ABSTRACT

Background: Human tissue/organ development is a complex, highly orchestrated process, regulated in part by the surrounding extracellular matrix (ECM). Every complex tissue, including the retina, has a unique ECM configuration that plays a critical role in cellular differentiation, adhesion, migration, and maturation. **Aim:** To characterize ECM expression of human induced pluripotent stem cell-derived optic vesicles (iPSC-OVs).

Methods: A 3-dimensional (3D) *in vitro* suspension culture system was used to direct differentiation of human induced pluripotent stem cells (iPSCs) into optic vesicles (OVs). Stepwise differentiation of iPSCs into retinal progenitor cells was confirmed by sequential expression of OTX2, SOX1, SIX6, LHX2, PAX6, and CHX10. Expression of ECM genes in iPSC-derived OV was analyzed by RT² Profiler™ PCR Array, whereas immunofluorescence staining was performed to detect ECM proteins in the OV.

Results: A number of cell adhesion molecules (CAMs) previously reported to be abundantly expressed in iPSCs such as E-cadherin, Intercellular adhesion molecule-1 (ICAM1), Integrin- α L, Integrin- α M, Integrin- α 6 were downregulated while neural and retina specific CAMs including neural cell adhesion molecule 1 (NCAM1), neural plakophilin-related armadillo repeat protein (NPRAP), Integrin- α 1 and Integrin- α 4 were upregulated. Several glycoproteins that have been reported to play key roles during retinogenesis, namely CD44, Tenascin C, Tenascin R, Neurocan, Neuroglycan C, Delta 2 Catenin, Vitronectin, and Reelin were also present.

Conclusion: We have identified an array of ECM proteins that were expressed during retinogenesis. Further characterization of these proteins will lead to a better understanding of retinal development.

© 2021 Heuy-Ching Wang. Hosting by Science Repository.

Introduction

Induced pluripotent stem cells (iPSCs) are derived from adult somatic cells that have been reprogrammed to become pluripotent. iPSCs share similar properties as embryonic stem cells (ESCs), including the plasticity to differentiate into any somatic cell [1, 2]. Several methods have been used to differentiate pluripotent stem cells into various cell types. One of the most common methods generates uniform and size-

controlled 3-dimensional (3D) stem cell aggregates called embryoid bodies (EBs) that are further differentiated into organoids of interest [3, 4]. Studies have shown that 3D EBs allow directed iPSC differentiation into desired cell types in a more efficient and controlled manner by recapitulating the key events that occur during early embryogenesis [5, 6]. EBs help to create a microenvironment that is conducive to cell-cell interactions and endogenous autocrine/paracrine signaling essential for self-renewal and differentiation.

*Correspondence to: Heuy-Ching Wang, Ph.D., Combat Casualty Care and Operational Medicine, Naval Medical Research Unit-San Antonio (NAMRU-SA), 3650 Chambers pass, Bldg. 3610, Fort Sam Houston, Texas 78234; Tel: 2105397017, E-mail: heuy-ching.h.wang.civ@mail.mil

Embryogenesis is a highly orchestrated process consisted of stepwise differentiation of pluripotent stem cells. During this process, embryos also start to secrete laminin and other extracellular matrix (ECM) molecules as early as the two-cell stage by a process called exocytosis [7]. As these embryonic cells further divide, each population of cells differentiates into tissue specific somatic cells. An EB-based differentiation approach has often been applied to induce *in vitro* retinogenesis of iPSCs, in which the sequential development of neuroectoderm specification, along with eye field and optic vesicle (OV) formation have been observed [8, 9]. ECM is a complex, dynamic network of proteins and carbohydrate molecules within the interstitial space and basement membrane of tissue [10, 11]. It is composed of different macromolecules including glycosylated proteins (proteoglycans), integrins, fibrous proteins and polysaccharides [10, 12, 13]. These molecules constitute the cellular microenvironment that provides structural support and exert influences on cell behaviors in various ways [14-17]. By binding to transmembrane proteins, ECM can regulate the underlying intracellular signaling pathways [18]. ECM can also regulate cell migration and differentiation by virtue of its stiffness and has been shown to influence tissue morphogenesis [19, 20]. In addition, ECM can sequester growth factors thereby influencing cell growth by regulating growth factor bioavailability.

ECM also plays a crucial role in degeneration/regeneration of tissue including retina in various diseases and injuries [21]. Several ECM molecules are upregulated in response to retinal damage. For example, Neuroglycan-C, a brain-specific chondroitin sulfate proteoglycan, has been shown to inhibit optic nerve regeneration after nerve crush injury in rats [22]. Other glycoproteins like Neurocan and CD44 are expressed at the outer interface of abutted retinas and inhibit neurite outgrowth [23]. Tenascin-C is also abundantly expressed after nerve injury which is important for axonal regeneration [24]. ECM remodeling also occurs in retinal diseases like glaucoma and diabetic retinopathy [25, 26].

In this study, we used a 3D *in vitro* suspension culture system to direct differentiation of human iPSCs into OVs and examined their ECM composition during early events of retinogenesis. Based on the results described here, we report that iPSCs undergoing retinal differentiation in 3D produce retina-specific ECMs. Further characterization of selected iPSC-derived ECM proteins can lead to a better understanding of retinal development and shed light on the potential application of stem cell-derived retinal organoids as a source of autologous ECM materials for retinal repair and regeneration.

Materials and Methods

I Culture and Maintenance of iPSCs

Human iPSC line IMR90-1 was obtained from WiCell (Madison, WI). iPSCs were seeded on Matrigel-coated (Corning, NY) 6 well-plates and maintained in mTeSR1 medium (Stem Cell Technologies, Vancouver, BC, Canada) at 37°C with 5% CO₂. Differentiated colonies were identified and removed via visual inspection using a dissecting microscope. Cells were manually passaged using Stempro EZPassage (ThermoFisher Scientific, Waltham, MA) when they reached 80% confluency, where approximately 40 to 50 colonies were seeded onto each well of the Matrigel-coated plates. iPSCs were tested for

pluripotency via immunostaining for various pluripotency markers as previously described [27].

II 3D *In Vitro* Retinal Differentiation

A retinal differentiation protocol was adapted from previous studies with modifications [9, 28]. Briefly, iPSCs were treated with 10 µM Rock inhibitor, Y-27632 (Stem Cell Technologies, Vancouver, Canada) for two hours, rinsed with Phosphate Buffered Saline (PBS), and dissociated into single cells using 0.1% Trypsin for 10 minutes. The cells were washed with PBS and resuspended at a density of 90,000 cells/mL in retinal differentiation medium containing G-MEM (ThermoFisher Scientific, Waltham, MA) with 20% KnockOut Serum Replacement (KSR) (ThermoFisher Scientific, Waltham, MA), 0.2 mM nonessential amino acids, 1 mM pyruvate, 0.1 mM 2-mercaptoethanol, 10 µM Y-27632, 3 µM IWR-1-endo (Wnt inhibitor, Calbiochem, Billerica, MA) and antibiotic/antimycotic (ThermoFisher Scientific). These cells were then assembled into uniform and size-controlled EBs of 9000 cells by aliquoting 100 µL of the cell suspension into each well of the Sumilon PrimeSurface 96-well V-bottomed plates (Sumitomo Bakelite, Novi, MI). After 24 hours, 1 µL of Matrigel at 1% v/v was added to each well. Half the medium in each well was changed to fresh medium without Y-27632 on Day 6. Cells were further cultured until Day 12.

On Day 12, the EBs were collected, washed twice and resuspended in medium containing G-MEM with 20% KSR, 10% Fetal bovine serum (FBS), 1% Matrigel, 0.2 mM nonessential amino acids, 1 mM pyruvate, 0.1 mM 2-mercaptoethanol, and 1X antibiotic/antimycotic cocktail and transferred to 6-well ultra-low attachment plate (Corning, Corning, NY) at a density of 30-40 EBs/well and cultured in suspension. Three µM CHIR 99021, (Wnt agonist, R&D Systems, Minneapolis, MN) and 100 nM smoothened agonist (SAG, R&D Systems, Minneapolis, MN) were added to the medium on Day 15. The medium was changed to neuroretina medium containing DMEM/F12-Glutamax, N2 supplement (ThermoFisher Scientific, Waltham, MA), and 1X antibiotic/antimycotic cocktail on Day 18. Cell differentiation was continued until Day 24.

III Immunofluorescence Staining of EBs During Stepwise Retinal Differentiation

EBs were harvested on Day 8, 12, and 24 during stepwise differentiation and fixed in 4% paraformaldehyde (PFA). After overnight fixation, the EBs were immersed in 15% and 30% sucrose (in PBS) for 2 hours each and embedded in optimal cutting temperature (OCT) compound. Embedded molds were frozen gradually and then sliced into 20 µm sections.

Cryosections were permeabilized with 0.1% Triton X-100 and incubated with antibodies against *OTX2*, *SOX1*, *SIX6*, *LHX2*, *PAX6*, or *CHX10* to detect markers of retinal development (Table 1). Fluorophore-conjugated secondary antibodies (Molecular Probes, Eugene, OR) were used for immunofluorescence detection. Cell nuclei were counterstained with 4, 6-diamidino-2-phenylindole (DAPI). Fluorescent images were acquired using a Leica TCS SP5 II confocal microscope (Leica Microsystems, Inc. Buffalo Grove, IL) and processed using the Image J software.

Table 1: Primary antibodies used for immunofluorescence assay.

Antibody	Type	Source	Dilution
CD44	Mouse monoclonal	Novus	1:100
<i>CHX10</i>	Rabbit polyclonal	Novus	1:100
<i>LHX2</i>	Goat polyclonal	Santa Cruz	1:25
Neurocan	Mouse monoclonal	ABCAM	1:50
Neuroglycan C	Rabbit polyclonal	ABCAM	1:300
<i>Pax6</i>	Mouse monoclonal	Developmental hybridoma bank	1:50
<i>Sox1</i>	Goat polyclonal	R&D Systems	1:40
<i>Six6</i>	Rabbit polyclonal	ABCAM	1:100
Tenascin C	Rabbit polyclonal	ABCAM	1:150
Tenascin R	Rabbit polyclonal	Novus	1:50

IV RT² Profiler™ PCR Array Assay

Total RNA was extracted from iPSCs (day 0) and iPSCs-derived OV_s on day 24 using the RNeasy mini kit (Qiagen, Valencia, CA). RNA was purified with RNase-free DNase (Qiagen). The purified RNA was reverse transcribed to cDNA using the RT First strand kit (Qiagen). cDNA was mixed with the RT SYBR Green ROX qPCR mastermix and aliquoted into the Human Extracellular Matrix and Cell Adhesion Molecules RT² Profiler™ PCR Array (Qiagen). Applied Biosystem 7300 RT PCR system (Applied Biosystems, Foster City, CA) was used for thermal cycling: segment 1 – 1 cycle: 95°C for 10 minutes, segment 2 – 40 cycles: 95°C for 15 seconds followed by 60°C for 1 minute. Data were analyzed using the RT² Profiler™ PCR Array Data Analysis Webportal (Link) according to the manufacturer's instructions. Fold changes in gene expression were calculated using the $\Delta\Delta C_t$ method. Expression of house-keeping genes was used to normalize the data and calculate $\Delta\Delta C_t$ values.

V Immunofluorescence Staining of ECM

EB_s on Day 24 post-differentiation were fixed and processed for immunofluorescence staining following the general protocol as described above. Antibodies against Neuroglycan C, Neurocan, Tenascin C, CD44, Tenascin R, Reelin, NCAM, Delta 2 Catenin, and Vitronectin were used to detect expression of ECM proteins. Fluorophore-conjugated secondary antibodies were used for immunofluorescence detection. DAPI was used to counterstain cell nuclei. Images were acquired using a confocal microscope as previously described. All images were processed using the Image J software.

VI Statistical Analysis

Each experiment was performed in triplicates. Data were analyzed by Student's t test and presented as the mean $\pm$ standard deviation (SD). A value of $p < 0.05$ was considered statistically significant.

Results

I In Vitro 3D Differentiation of OV_s

We applied a well-established retinal differentiation protocol to direct differentiation of iPSCs into OV_s with some modifications [9, 28]. IMR90-1 cells were dissociated into single cells and aggregated to form uniform and size-controlled EB_s of 9000 cells/EB (Figure 1A). We observed that IMR90-1 cells underwent stepwise differentiation into retinal progenitors. They expressed neuroectodermal marker *OTX2* and *SOX1* by Day 8 (Figure 1B) and acquired eye field specification markers, *SIX6* and *LHX2* by Day 12 (Figure 2). These EB_s further developed into OV_s as marked by co-expression of neuroretina markers *PAX6* and *CHX10* (Figure 3). Here we have shown that 3D *in vitro* differentiation of iPSCs into OV_s recapitulates *in vivo* retinogenesis via sequential expression of the neuroectoderm, eye field and neuroretina markers.

II Expression Profile of ECM Genes in iPSC-Derived OV_s

Figures 3 showed that OV_s derived from iPSCs in 3D culture expressed markers of the neuroretina. These results suggest that iPSC-derived OV_s can be used as a model to study the expression of retina-specific ECM. To characterize the ECM, gene expression profiles of iPSCs and iPSC-derived OV_s were analyzed with a commercially available RT² Profiler™ PCR Array. iPSCs-derived OV_s showed significantly downregulated expression of CAMs including E-cadherin, Intercellular adhesion molecule-1 (ICAM1), Integrin- α L, Integrin- α M, and Integrin- α 6 (Figure 4) and upregulated expression of Neural cell adhesion

molecule 1 (NCAM1), Neural plakophilin related armadillo protein (NPRAP), Integrin- α 1 and Integrin- α 4 (Figure 4). Furthermore, ECM expression of glycoproteins Tenascin C, CD44, Vitronectin and

Fibronectin I was significantly upregulated in iPSCs-derived OV's (Figure 5).




Figure 1: Stepwise differentiation of iPSCs towards a retinal lineage. **A)** 9000 iPSCs were aggregated to form uniform-sized EBs and differentiated towards a retinal lineage. **B)** Differentiating EBs expressed neuronal marker *OTX2* and *SOX1* by Day 8 (scale bar=200 μ m).




Figure 2: Differentiating EBs expressed eye field specification markers *SIX6* and *LHX2* by Day 12 (scale bar= 200 μ m).




Figure 3: Optic vesicle formation from iPSCs. Developing EBs from iPSCs co-expressed neuronal markers *PAX6* and retina progenitor cell marker *CHX10* (scale bar=25µm).




Figure 4: Expression of CAMs and glycoproteins in iPSC-derived OVVs on Day 24 of differentiation. iPSC-specific CAMs including E-cadherin, ICAM1, Integrin-α L, Integrin-α M, Integrin-α 6 were downregulated while neural and retina specific CAMs NCAM1, NPRAP, Integrin-α 1 and Integrin-α 4 were upregulated.




Figure 5: Expression of glycoproteins in iPSC-derived OVVs on Day 24 of differentiation. Genes for ECM proteins important for retinogenesis namely Tenascin-C, CD44, Fibronectin I and Vitronectin were upregulated.

III Expression of ECM Proteins in iPSC-Derived OV_s

The data from the RT² ProfilerTM PCR array showed that iPSCs-derived OV_s expressed several ECM genes associated with cells of the retinal lineage. Immunofluorescence analysis was performed to further verify the expression and localization of ECM proteins. CD44, a glycoprotein expressed in retinal Müller cells was significantly higher in iPSC-

derived OV_s as detected by RT-PCR microarray (Figure 4) and via immunofluorescence staining (Figure 7) [29]. Tenascin R, a known modulator of retinal axon growth, was also detected in iPSC-derived OV_s (Figure 8) [30]. Additional neural and retinal ECM proteins were found to be expressed in the iPSC-derived OV_s, include Reelin, Vitronectin, Delta 2 Catenin, Neuroglycan C, Neurocan, and Tenascin C (Figures 6-8) [31-33].




Figure 6: Protein expression of Neuroglycan C, Neurocan, and Tenascin C in iPSC-derived OV_s. Neuroglycan C, Neurocan, and Tenascin C were abundantly expressed in the developing iPSC-derived OV_s (scale bar=25μm). DAPI (middle column), merge view (right column). These ECM proteins are known to be expressed in the developing eye *in vivo*.




Figure 7: Expression of NCAM, CD44 and Vitronectin in iPSCs-derived OV_s. Immunofluorescence analysis showed that OV_s expressed NCAM, CD 44 and Vitronectin in the developing OV_s (left column). Nuclear staining/DAPI (middle column), merge view (right column), scale bar=50μm.




Figure 8: Expression of Reelin, Tenascin R and Delta 2 Catenin in iPSC-derived OVs. Retina abundant proteins like Reelin, Tenascin-R and Delta 2 Catenin were expressed in developing iPSC-derived OVs during early stage retinogenesis. Nuclear staining/DAPI (middle column), merge view (right column), scale bar=50 μ m.

Discussion

Intracellular and extracellular changes occur during tissue/organ development and provide necessary cues for differentiation to ensure cells can acquire a tissue-specific phenotype [17, 19, 27, 34]. Engineering iPSCs into 3D EBs provides a unique environment that mimics the environment of early embryogenesis [29, 30]. Pluripotent stem cells have shown stepwise acquisition of developmental phenotypes during differentiation towards a defined lineage [3]. During human retinogenesis, the appearance of eye field cells (*SIX6*⁺ and *LHX2*⁺) in the diencephalon of the anterior neuroectoderm (with *SOX1*⁺ and *OTX2*⁺ cells) marks the initiation of the neuroretina development (with *CHX10*⁺ and *PAX6*⁺ retinal progenitor cells). The formation of OVs followed by subsequent invagination of the vesicles results in morphogenesis of the optic cup [8, 31, 32]. Similar sequences of retinal morphogenesis were reported in 3D differentiation of human ESCs and iPSCs *in vitro* [9, 28]. Even though cellular and phenotypical changes that occur during specification towards a retinal lineage have been well-documented in literature, little is known about the dynamics of ECM formation during retinal development.

In this study, we have demonstrated that iPSC-derived OVs expressed several CAMs/ECMs that have previously been shown to play key roles in retinogenesis *in vivo*, specifically NCAM1, NPRAP, Tenascin C, CD44, Fibronectin I, and Vitronectin. The iPSCs-derived OVs also expressed Tenascin R, Neurocan, Neuroglycan C, Reelin, Laminin S, and Delta 2 Catenin. Neuroglycan C is a neural tissue specific transmembrane chondroitin sulfate proteoglycan that regulates the formation of the retinal neural network [35, 36]. Another glycoprotein,

Tenascin C works in conjunction with Tenascin R to regulate axonal growth during retinogenesis [24, 37]. Furthermore, Tenascin C has also been shown to interact with Neurocan, another glycoprotein that is expressed in neuronal tissue [38, 39]. CD44, another cell surface glycoprotein that was upregulated in this study, is known to be expressed by Müller cells in the retina *in vivo* [40, 41]. Finally, Fibronectin, and Vitronectin are integral components of the human retina which were also found to be significantly overexpressed in OVs derived from iPSCs [38-41].

The application of biological ECM scaffolds for tissue repair and regeneration is a rapidly advancing interdisciplinary field [33, 42]. A recent study showed that decellularized bovine retinal ECM can support attachment and growth of human retinal progenitor cells [43]. ECM scaffolds derived from CNS tissue have also been reported to support migration and proliferation of neural stem cells [44, 45]. Furthermore, another study demonstrated that ECM scaffolds derived from skin promoted cell engraftment, proliferation and wound healing without scar formation [46, 47]. Many groups have confirmed that tissue-specific ECM promotes maturation and integration when seeded with corresponding cell types [7, 20, 48-54]. These studies indicate that native ECM is a good candidate as a biomaterial for tissue engineering applications.

ECM can be acquired from different sources including human (allogeneic) or animal (xenogeneic) donor tissues [48]. However, it is important to note that animal-derived ECM products pose risks of immunoreactivity and zoonotic disease transmission, whereas human donor tissues are often in short supply [50, 55-60]. Since patient-derived

materials are generally preferred for tissue replacement therapy, generating ECM from iPSC-derived organoids could provide autologous materials ideal for tissue repair and regeneration. Using stem cells as a potential source for tissue-specific ECM is a novel approach. There are a few reports on stem cell-derived ECM applications in the literature, including a recent study that described the ability of ECM derived from undifferentiated EBs to support ESC proliferation and differentiation [7]. Several studies have shown that while undergoing chondrogenic differentiation, MSC-derived ECM can induce chondrogenic differentiation of seeded stem cells without exogenous growth factors [61, 62]. These studies have demonstrated the feasibility of obtaining chondrogenic ECM for cartilage regeneration from MSC aggregates differentiating towards a chondrogenic phenotype.

In summary, we have successfully used human iPSCs as a tool to recapitulate early stages of retinal development in this study. We have shown that developing OV's expressed several ECM proteins that are known to play critical roles during retinogenesis *in vitro*. We have also demonstrated that iPSCs in 3D suspension culture can generate native retina-specific ECM. Future work will include examining the potential of iPSC-derived OV's as an autologous source of ECM materials that can support axonal survival, growth, and guidance for retinal regeneration. Further studies will also be needed to validate the regenerative potency of autologous retina-specific ECM in retinal disease/injury.

Acknowledgement

This work was supported by the U.S. Army Clinical Rehabilitative Medicine Research Program (CRM RP), Military Operational Medicine Research Program (MOM RP) and Oak Ridge Institute for Science and Education (ORISE).

Disclaimers

The opinions or assertions contained herein are the private views of the authors and are not to be construed as official or as reflecting the views of the Department of the Army or the Department of Defense.

Author Contributions

Kaini RR performed data collection, analysis and wrote the manuscript. Rettinger CL performed data analysis and wrote the manuscript. Wang HC and Kaini RR designed the experiments. Wang HC obtained funding, supervised the study, and revised the manuscript. All authors approved the manuscript.

REFERENCES

- Robinton DA, Daley GQ (2012) The promise of induced pluripotent stem cells in research and therapy. *Nature* 481: 295-305. [Crossref]
- Yu J, Vodyanik MA, Smuga Otto K, Antosiewicz Bourget J, Frane JL et al. (2007) Induced pluripotent stem cell lines derived from human somatic cells. *Science* 318: 1917-1920. [Crossref]
- Ader M, Tanaka EM (2014) Modeling human development in 3D culture. *Curr Opin Cell Biol* 31: 23-28. [Crossref]
- Clevers H (2016) Modeling Development and Disease with Organoids. *Cell* 165: 1586-1597. [Crossref]
- Bratt Leal AM, Carpenedo RL, McDevitt TC (2009) Engineering the embryoid body microenvironment to direct embryonic stem cell differentiation. *Biotechnol Prog* 25: 43-51. [Crossref]
- Yodmuang S, Marolt D, Marcos Campos I, Gadjanski I, Vunjak Novakovic G et al. (2015) Synergistic effects of hypoxia and morphogenetic factors on early chondrogenic commitment of human embryonic stem cells in embryoid body culture. *Stem Cell Rev Rep* 11: 228-241. [Crossref]
- Goh SK, Olsen, Banerjee I (2013) Extracellular matrix aggregates from differentiating embryoid bodies as a scaffold to support ESC proliferation and differentiation. *PLoS One* 8: e61856. [Crossref]
- Meyer JS, Howden SE, Wallace KA, Verhoeven AD, Wright LS et al. (2011) Optic vesicle-like structures derived from human pluripotent stem cells facilitate a customized approach to retinal disease treatment. *Stem Cells* 29: 1206-1218. [Crossref]
- Nakano T, Ando S, Takata N, Kawada M, Muguruma K et al. (2012) Self-formation of optic cups and storable stratified neural retina from human ESCs. *Cell Stem Cell* 10: 771-785. [Crossref]
- Frantz C, Stewart KM, Weaver VM (2010) The extracellular matrix at a glance. *J Cell Sci* 123: 4195-4200. [Crossref]
- Tanzer ML (1985) Extracellular matrix: extracellular matrix biochemistry. *Science* 227: 289-290. [Crossref]
- Halper J, Kjaer M (2014) Basic components of connective tissues and extracellular matrix: elastin, fibrillin, fibulins, fibrinogen, fibronectin, laminin, tenascins and thrombospondins. *Adv Exp Med Biol* 802: 31-47. [Crossref]
- Kular JK, Basu S, Sharma RI (2014) The extracellular matrix: Structure, composition, age-related differences, tools for analysis and applications for tissue engineering. *J Tissue Eng* 5: 2041731414557112. [Crossref]
- Boretti MI, Gooch KJ (2008) Effect of extracellular matrix and 3D morphogenesis on islet hormone gene expression by Ngn3-infected mouse pancreatic ductal epithelial cells. *Tissue Eng Part A* 14: 1927-1937. [Crossref]
- Faulk DM, Johnson SA, Zhang L, Badyak SF (2014) Role of the extracellular matrix in whole organ engineering. *J Cell Physiol* 229: 984-989. [Crossref]
- Streuli C (1999) Extracellular matrix remodelling and cellular differentiation. *Curr Opin Cell Biol* 11: 634-640. [Crossref]
- Watt FM, Huck WT (2013) Role of the extracellular matrix in regulating stem cell fate. *Nat Rev Mol Cell Biol* 14: 467-473. [Crossref]
- Vecino E, Heller JP, Veiga Crespo P, Martin KR, Fawcett JW (2015) Influence of extracellular matrix components on the expression of integrins and regeneration of adult retinal ganglion cells. *PLoS One* 10: e0125250. [Crossref]
- Bonnans C, Chou J, Werb Z (2014) Remodelling the extracellular matrix in development and disease. *Nat Rev Mol Cell Biol* 15: 786-801. [Crossref]
- Varshney S, Hunter DD, Brunken WJ (2015) Extracellular Matrix Components Regulate Cellular Polarity and Tissue Structure in the Developing and Mature Retina. *J Ophthalmic Vis Res* 10: 329-339. [Crossref]
- Ren T, van der Merwe Y, Steketee MB (2015) Developing Extracellular Matrix Technology to Treat Retinal or Optic Nerve Injury(1,2,3). *eNeuro* 2: ENEURO.0077-15.2015. [Crossref]
- Reinhard J, Renner M, Wiemann S, Shakoor DA, Stute G et al. (2017) Ischemic injury leads to extracellular matrix alterations in retina and optic nerve. *Sci Rep* 7: 43470. [Crossref]

23. Zhang Y, Kardaszewska AK, van Veen T, Rauch U, Perez MTR et al. (2004) Integration between abutting retinas: role of glial structures and associated molecules at the interface. *Invest Ophthalmol Vis Sci* 45: 4440-4449. [\[Crossref\]](#)
24. Besser M, Jagatheaswaran M, Reinhard J, Schaffelke P, Faissner A (2012) Tenascin C regulates proliferation and differentiation processes during embryonic retinogenesis and modulates the de-differentiation capacity of Müller glia by influencing growth factor responsiveness and the extracellular matrix compartment. *Dev Biol* 369: 163-176. [\[Crossref\]](#)
25. Roy S, Amin S, Roy S (2016) Retinal fibrosis in diabetic retinopathy. *Exp Eye Res* 142: 71-75. [\[Crossref\]](#)
26. Schneider M, Fuchshofer R (2016) The role of astrocytes in optic nerve head fibrosis in glaucoma. *Exp Eye Res* 142: 49-55. [\[Crossref\]](#)
27. Kaini RR, Shen Gunther J, Cleland JM, Greene WA, Wang HC et al. (2016) Recombinant Xeno-Free Vitronectin Supports Self-Renewal and Pluripotency in Protein-Induced Pluripotent Stem Cells. *Tissue Eng Part C Methods*. [\[Crossref\]](#)
28. Sasai Y, Eiraku M, Suga H (2012) In vitro organogenesis in three dimensions: self-organising stem cells. *Development* 139: 4111-4121. [\[Crossref\]](#)
29. Sutha K, Schwartz Z, Wang Y, Hyzy S, Boyan BD et al. (2015) Osteogenic embryoid body-derived material induces bone formation in vivo. *Sci Rep* 5: 9960. [\[Crossref\]](#)
30. Sepulveda DE, Andrews BA, Asenjo JA, Papoutsakis ET (2008) Comparative transcriptional analysis of embryoid body versus two-dimensional differentiation of murine embryonic stem cells. *Tissue Eng Part A* 14: 1603-1614. [\[Crossref\]](#)
31. Chow RL, Lang RA (2001) Early eye development in vertebrates. *Annu Rev Cell Dev Biol* 17: 255-296. [\[Crossref\]](#)
32. Eiraku M, Takata N, Ishibashi H, Kawada M, Sakakura E et al. (2011) Self-organizing optic-cup morphogenesis in three-dimensional culture. *Nature* 472: 51-56. [\[Crossref\]](#)
33. O'Brien FJ (2011) Biomaterials & scaffolds for tissue engineering. *materialstoday* 14: 88-95.
34. Hausman RE (2007) Ocular extracellular matrices in development. *Prog Retin Eye Res* 26: 162-188. [\[Crossref\]](#)
35. Escher P, Cottet S, Aono S, Oohira A, Schorderet DF (2008) Differential neuroglycan C expression during retinal degeneration in Rpe65^{-/-} mice. *Mol Vis* 14: 2126-2135. [\[Crossref\]](#)
36. Inatani M, Tanihara H, Oohira A, Otori Y, Nishida A et al. (2000) Neuroglycan C, a neural tissue-specific transmembrane chondroitin sulfate proteoglycan, in retinal neural network formation. *Invest Ophthalmol Vis Sci* 41: 4338-4346. [\[Crossref\]](#)
37. Schumacher S, Stube EM (2003) Regulated binding of the fibrinogen-like domains of tenascin-R and tenascin-C to the neural EGF family member CALEB. *J Neurochem* 87: 1213-1223. [\[Crossref\]](#)
38. Inatani M, Tanihara H (2002) Proteoglycans in retina. *Prog Retin Eye Res* 21: 429-447. [\[Crossref\]](#)
39. Kwan KM (2014) Coming into focus: the role of extracellular matrix in vertebrate optic cup morphogenesis. *Dev Dyn* 243: 1242-1248. [\[Crossref\]](#)
40. Chaitin MH, Ankrum MT, Wortham HS (1996) Distribution of CD44 in the retina during development and the rds degeneration. *Brain Res Dev Brain Res* 94: 92-98. [\[Crossref\]](#)
41. Reinhard J, Joachim SC, Faissner A (2015) Extracellular matrix remodeling during retinal development. *Exp Eye Res* 133: 132-140. [\[Crossref\]](#)
42. Yi S, Ding F, Gong L, Gu X (2017) Extracellular Matrix Scaffolds for Tissue Engineering and Regenerative Medicine. *Curr Stem Cell Res Ther* 12: 233-246. [\[Crossref\]](#)
43. Kundu J, Michaelson A, Talbot K, Baranov P, Young MJ et al. (2016) Decellularized retinal matrix: Natural platforms for human retinal progenitor cell culture. *Acta Biomater* 31: 61-70. [\[Crossref\]](#)
44. Crapo PM, Medberry CJ, Reing JE, Tottey S, van der Merwe Y et al. (2012) Biologic scaffolds composed of central nervous system extracellular matrix. *Biomaterials* 33: 3539-3547. [\[Crossref\]](#)
45. Crapo PM, Tottey S, Slivka PF, Badyak SF (2014) Effects of biologic scaffolds on human stem cells and implications for CNS tissue engineering. *Tissue Eng Part A* 20: 313-323. [\[Crossref\]](#)
46. Wainwright DJ (1995) Use of an acellular allograft dermal matrix (AlloDerm) in the management of full-thickness burns. *Burns* 21: 243-248. [\[Crossref\]](#)
47. Zhong SP, Zhang YZ, Lim CT (2010) Tissue scaffolds for skin wound healing and dermal reconstruction. *Wiley Interdiscip Rev Nanomed Nanobiotechnol* 2: 510-255. [\[Crossref\]](#)
48. Choi YC, Choi JS, Kim BS, Kim JD, Yoon HI et al. (2012) Decellularized extracellular matrix derived from porcine adipose tissue as a xenogeneic biomaterial for tissue engineering. *Tissue Eng Part C Methods* 18: 866-876. [\[Crossref\]](#)
49. Cigognini D, Lomas A, Kumar P, Satyam A, English A et al. (2013) Engineering in vitro microenvironments for cell based therapies and drug discovery. *Drug Discov Today* 18: 1099-1108. [\[Crossref\]](#)
50. Eitan Y, Sarig U, Dahan N, Machluf M (2010) Acellular cardiac extracellular matrix as a scaffold for tissue engineering: in vitro cell support, remodeling, and biocompatibility. *Tissue Eng Part C Methods* 16: 671-683. [\[Crossref\]](#)
51. Lancaster MA, Knoblich JA (2014) Organogenesis in a dish: modeling development and disease using organoid technologies. *Science* 345: 1247125. [\[Crossref\]](#)
52. Lin YM, Zhang A, Rippon HJ, Bismarck A, Bishop AE (2010) Tissue engineering of lung: the effect of extracellular matrix on the differentiation of embryonic stem cells to pneumocytes. *Tissue Eng Part A* 16: 1515-1526. [\[Crossref\]](#)
53. Turner AEB, Flynn LE (2012) Design and characterization of tissue-specific extracellular matrix-derived microcarriers. *Tissue Eng Part C Methods* 18: 186-197. [\[Crossref\]](#)
54. Xing Q, Yates K, Tahtinen M, Shearier E, Qian Z et al. (2015) Decellularization of fibroblast cell sheets for natural extracellular matrix scaffold preparation. *Tissue Eng Part C Methods* 21: 77-87. [\[Crossref\]](#)
55. Denner J, Mueller NJ (2015) Preventing transfer of infectious agents. *Int J Surg* 23: 306-311. [\[Crossref\]](#)
56. Tang C, Jin C, Du X, Yan C, Min BH et al. (2014) An autologous bone marrow mesenchymal stem cell-derived extracellular matrix scaffold applied with bone marrow stimulation for cartilage repair. *Tissue Eng Part A* 20: 2455-2462. [\[Crossref\]](#)
57. Turner NJ, Yates AJ Jr, Weber DJ, Qureshi IR, Stolz DB et al. (2010) Xenogeneic extracellular matrix as an inductive scaffold for regeneration of a functioning musculotendinous junction. *Tissue Eng Part A* 16: 3309-3317. [\[Crossref\]](#)
58. Unger C, Skottman H, Blomberg P, Dilber MS, Hovatta O et al. (2008) Good manufacturing practice and clinical-grade human embryonic stem cell lines. *Hum Mol Genet* 17: R48-R53. [\[Crossref\]](#)
59. Fong AH, Romero López M, Heylman CM, Keating M, Tran D et al. (2016) Three-dimensional adult cardiac extracellular matrix promotes

- maturation of human induced pluripotent stem cell-derived cardiomyocytes. *Tissue Eng Part A* 22: 1016-1025. [[Crossref](#)]
60. Pati F, Jang J, Ha DH, Kim SW, Rhie JW et al. (2014) Printing three-dimensional tissue analogues with decellularized extracellular matrix bioink. *Nat Commun* 5: 3935. [[Crossref](#)]
61. Cai R, Nakamoto T, Kawazoe N, Chen G (2015) Influence of stepwise chondrogenesis-mimicking 3D extracellular matrix on chondrogenic differentiation of mesenchymal stem cells. *Biomaterials* 52: 199-207. [[Crossref](#)]
62. Mehlhorn AT, Niemeyer P, Kaiser S, Finkenzeller G, Stark GB et al. (2006) Differential expression pattern of extracellular matrix molecules during chondrogenesis of mesenchymal stem cells from bone marrow and adipose tissue. *Tissue Eng* 12: 2853-2862. [[Crossref](#)]